

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|---|-----------|--|
| (51) International Patent Classification ⁶: G01N 33/53, 33/68 | A1 | (11) International Publication Number: WO 00/02046 (43) International Publication Date: 13 January 2000 (13.01.00) |
| (21) International Application Number: PCT/SE99/01208 (22) International Filing Date: 2 July 1999 (02.07.99) (30) Priority Data: 9802402-9 3 July 1998 (03.07.98) SE 60/091,741 6 July 1998 (06.07.98) US (71) Applicant (for all designated States except US): KAROLIN- SKA INNOVATIONS AB [SE/SE]; S-171 77 Stockholm (SE). (72) Inventor; and (75) Inventor/Applicant (for US only): FROSTEGÅRD, Johan [SE/SE]; Törnrosavägen 9, S-131 47 Nacka (SE). (74) Agent: DR. LUDWIG BRANN PATENTBYRÅ AB; P.O. Box 1344, S-751 43 Uppsala (SE). | | (81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BI, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> |
| (54) Title: METHOD OF DIAGNOSING CARDIOVASCULAR DISEASE AND EARLY ATHEROSCLEROSIS | | |
| (57) Abstract | | |
| <p>The present invention provides a method of diagnosing cardiovascular disease and increased risk of early atherosclerosis in a human wherein the presence and/or concentration of antibodies to platelet activating factor (PAF) in a sample of body fluid of said human is assessed. The present invention further provides a kit carrying out said method.</p> | | |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|---------------------------------------|----|---|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

METHOD OF DIAGNOSING CARDIOVASCULAR DISEASE AND EARLY ATHEROSCLEROSIS

FIELD OF THE INVENTION

The present invention relates generally to methods for identifying patients who have cardiovascular disease and increased risk of developing atherosclerosis. More particularly, the invention relates to the detection of IgG antibodies to platelet activating factor (PAF) in body fluids of patients. The present inventors have shown that elevated concentrations of antibodies to PAF in body fluids is correlated to borderline hypertension and metabolic syndrome, i.e. early cardiovascular disease, which is connected to increased risk of developing early atherosclerosis.

BACKGROUND OF THE INVENTION

The morbidity and mortality associated with cardiovascular diseases and atherosclerosis in developed countries is higher than that associated with any other disorder. Hypertension is, together with hyperlipidemia, the most prominent risk factor for atherosclerosis. Individuals with borderline hypertension are an example of early cardiovascular disease in general, with endothelial dysfunction and increased risk of atherosclerotic disease, in apparently healthy individuals. Early atherosclerosis manifests itself in the form of cholesterol depositions in the arterial wall. During recent years, it has been convincingly shown that the atherosclerotic process is a chronic inflammation, characterized by presence of activated T cells and monocytes/macrophages. Many of these macrophages have developed into cholesterol-filled foam cells. The deposition is slow and starts at an early age. Clinical symptoms may take years to manifest themselves and are very serious; they include coronary heart disease and stroke. Generally, the disease process will have begun long before these clinical manifestations appear. There are available a number of genetic analysis screening for patients with pre-deposition for atherosclerosis. But it is desirable to have available a diagnostic technique which provides an early warning of the onset of the deposition. The importance of early detection is stressed by the fact that an effective long-term treatment is possible. The present techniques for diagnosing atherosclerosis depend on measuring cholesterol or triglycerid levels in serum or detection of atheromatous lesions, but by the time of detection, the most effective time for treatment has been passed. U.S. patent 5,731,208 discloses a screening test for atherosclerosis comprising determining the concentration of p-hydroxyphenylaldehyde-lysine in serum or plasma.

The present inventors have found that elevated concentrations of IgG antibodies to platelet activating factor (PAF) in patients are an indicator of cardiovascular diseases which is often accompanied by early atherosclerosis. More specifically, antibodies to PAF (aPAF) are associated with early vascular disease in the form of both borderline hypertension and the metabolic syndrome, both of which are strong risk factors for later stages of atherosclerosis, which give rise to clinical symptoms.

These results demonstrate that antibodies against PAF represent a novel category of anti-phospholipid antibodies (aPL), which are sensitive to early vascular dysfunction and disease, especially early atherosclerosis and hypertension.

aPL in general, especially against cardiolipin have been shown to predict risk of cardiovascular disease, also in autoimmune diseases like systemic lupus erythematosus (SLE) and our data thus indicate that antibodies against PAF is a novel category of aPL, with a potential as a marker also in other autoimmune conditions in addition to cardiovascular disease and atherosclerosis in general. aPL have been related to both arterial and venous thrombosis, and also to spontaneous abortion. These data indicate that antibodies to PAF were much more strongly associated with spontaneous abortion than aPL, and furthermore, that antibodies to PAF was a novel marker for disease activity in SLE.

Antibodies to PAF are therefore relevant also in these other autoimmune vascular-related diseases.

Also antibodies to PAF-like lipids are relevant in this context, one being lysophosphatidylcholine, where the results indicate a comparable profile as the one obtained by PAF antibodies.

Accordingly, it is a principal object of the present invention to provide a diagnostic method or screening test for early atherosclerosis or cardiovascular changes related to early atherosclerosis. It is yet another object of the invention to provide a kit for assaying the concentrations of aPAF for diagnosing early atherosclerosis or cardiovascular changes.

"Early atherosclerosis" as used herein refers to the very first stage of atherosclerosis, before the clinical onset of symptoms. "Early cardiovascular disease" as used herein refers to the first stages of cardiovascular disease, as in

borderline hypertension and the metabolic syndrome, when atherosclerosis is yet not easy to detect by other methods and has not given rise to disease.

Platelet activating factor (PAF) is a phospholipid inflammatory mediator that is synthesized by a variety of cells, including monocytes and endothelial cells. During oxidation of LDL, PAF-like lipids are produced. PAF may therefore be of importance in pathological processes in the vascular wall like atherosclerosis and hypertension. In a previous report, the existence of antibodies to PAF (aPAF) were described in individuals with phospholipid antibody syndrome (Barquinero et al., 1994.), but nothing has been reported about possible clinical implications of these antibodies and a putative role in cardiovascular disease.

DISCLOSURE OF THE INVENTION

As mentioned above, we have surprisingly shown that concentration of antibodies to PAF (aPAF) is an effective indicator of early cardiovascular disease. We have found that antibodies to this particular antigen develop in patients well before the clinical onset of atherosclerosis.

In our study we found that concentration of aPAF was 49.3% higher in borderline hypertension men than in normotensive men. When defining aPAF concentrations above mean concentration of control plus two standard deviations (i.e. $0.144 + (2 \times 0.109) = 0.362$ OD405) as positive, 15 men out of 73 were positive in the borderline hypertension group whereas only 3 men out of 73 were positive in the normotensive group. Antibodies to PAF as a marker for early atherosclerosis may be combined with additional and alternative markers for early atherosclerosis to improve the accuracy of the diagnosis, such as determining the concentrations of cholesterol, blood lipids or p-hydroxyphenylaldehyde-lysine.

Antibodies against PAF (aPAF) may be determined using any of the methods and techniques conventional in the art for such determination. Conveniently, such a method may comprise immunoassay e.g. ELISA or RIA. The immunoassay will conveniently use an antigen (PAF) in immobilized form, e.g. on microtitre plates, membranes or beads, to isolate the target aPAF. In a sandwich assay, the bound antigen may be labelled using additional soluble antibody, which may be monoclonal or polyclonal and which may either carry a label or, more conveniently, may itself be labelled subsequently by reaction with a secondary antibody carrying a label. Suitable labels include radionuclides, fluorescent substances, and enzymes.

Alternatively, a competitive binding assay may be used. Conveniently, the components needed to perform the immunoassay will be supplied in kit form. Such a kit would comprise:

- a) an antigen capable of binding to aPAF and, optionally;
- a labelled sample of antigen to aPAF or a fragment thereof;
- said antigen (a) in non-immobilised form;
- a labelled secondary antibody specific to said antigen (c).

The body fluid on which the determination is performed may be any body fluid in which aPAF may be located, but conveniently will be or serum or plasma. In some cases it may be convenient to extract the antibodies, or otherwise treat the sample prior to determination.

The invention will now be described in greater detail by reference to the following non-limiting examples:

EXAMPLE 1

Determination of concentration of PAF antibodies of early atherosclerosis patients and of normal patients.

In order to investigate the role of aPAF in borderline hypertension (BHT) and early atherosclerosis, we studied a group of 146 middle aged men, where borderline hypertension were compared with age-matched controls. We here report that serum aPAF titers are enhanced in patients with borderline hypertension and metabolic syndrome.

Patients were recruited from a population screening program as previously described (Lemne et al 1995). BHT was defined as diastolic blood pressure (DBP) of 85 to 94 mmHg, and the screening identified 81 men who remained within the range for borderline hypertension during repeated measurements over a three year period. From the same population 80 age matched controls were recruited, whose blood pressure was measured on two occasions a few weeks apart, and was <80 mmHg on both occasions.

Of the 81 men with BHT and the 80 NT controls who agreed to participate, 73 in the BHT and 75 in the NT group completed all procedures of the present study.

None of the subjects had any other illnesses or were regularly using any drugs known to influence blood pressure, metabolic or inflammatory variables.

All subjects were investigated according to the same schedule. Both BHT and NT controls were investigated simultaneously when possible and no more than 4 weeks apart. Blood samples for analyses of metabolic and inflammatory variables were taken between 8 and 9:30 a.m., after 8 to 12 hours of fasting. All samples were drawn after 15 minutes of rest in the supine position.

An identical procedure was followed at each occasion during the entire recruitment period. All blood pressure measurements were performed with a mercury sphygmomanometer. The cuff was adjusted according to the circumference of the arm and placed at the level of the heart. Blood pressure was recorded as the mean of two measurements taken after 5 minutes rest in the supine position. Systolic and diastolic blood pressure measurements were defined according to Korotkoff I and V. The same specially trained nurse performed the measurements on all occasions.

The right and left carotid arteries were examined with a duplex scanner (Acuson 128XP/5, Mountain View, CA, USA) using a 7.0 MHz linear array transducer. The subjects were investigated in the supine position and intima-media (I-M) thickness was determined in the far wall as the distance between the leading edge of the lumen-intima echo and the leading edge of the media-adventitia's echo. Plaque was defined as a localized I-M thickening with a thickness > 1mm and a 100% increase in thickness compared with normal, adjacent wall segments. Plaque occurrence was scored as present or absent. Plaque was screened for in the common, internal and external carotid arteries on both sides, as described earlier (Lemne et al 1995).

All patients were weighed without other clothing than underwear, using the same scale (Delta 707, SECA, Germany). Length was measured with a special ruler, fixed to the wall. Waist circumference was measured at the level of the umbilicus and the hips were measured at the level of the greatest circumference. Body mass index (BMI) was subsequently calculated as weight in kilograms/(height in meters)².

IgG antibodies to PAF were determined according to Example 2.

PAF (1-O-alkyl-2-acetyl-sn-glycero-3-phosphocholine) was obtained from Sigma, St Louis, USA.

Lipid and lipoprotein levels were determined by a combination of preparative ultracentrifugation followed by lipid analyses in the lipoprotein fractions as previously described (Lemne et al 1995).

Venous blood samples for determination of plasma insulin (Radio-Immuno Assay, Kabi Pharmacia, Sweden) were taken.

Serum immunoglobulins, IgG, IgM and IgA were determined as described (Frostegård et al, Hypertension 1997)

Variables were tested for skewness. For skewed variables non-parametric tests were used for comparisons between the groups (Mann-Whitney U-test), whereas Student's t-test was used for normally distributed variables. Spearman rank correlation coefficients were calculated to estimate interrelations between antibody levels, metabolic variables and blood pressure levels. The significance level was put at $p < 0.05$. Values in the text are given as mean \pm standard deviation (SD) as indicated.

Results

Basic characteristics of the two study groups are presented in Table 1. The mean blood pressure level in the NT group was 125/75 ($\pm 11/\pm 5$) mmHg as compared to 141/89 ($\pm 10/\pm 2$) mmHg in the BHT group. The two groups were well-matched for age. The BHT men had a significantly altered metabolic profile with fasting hyperinsulineamia and dyslipoproteineamia, as previously presented (Table I). In the BHT group 26 % of the subjects had plaque on one or both sides while and the corresponding figure for the NT group was 16% (19 vs 10 subjects, n.s.).

In the material as a whole, the aPAF levels were significantly higher in the BHT group, compared with the NT group (Table II). There was no difference in alysoPAF levels between the BHT and NT group.

There were no significant differences in aPAF levels between individuals with plaque (n=29) compared to individuals without (n=117); data not shown).

If values above 2SD in the control group were defined as positive, 21% in the BHT group and 4% in the NT group had increased aPAF levels. Age did not correlate with antibody levels (data not shown).

To exclude the possibility that differences in antibody levels simply reflected enhanced total antibody levels total IgG was determined. There was no difference between the BHT group and controls (data not shown).

In the material as a whole, and the two groups separately there were no significant correlations between aPAF and BMI, blood pressure levels, or smoking (data not shown).

However, individuals with the metabolic syndrome (defined as having at least two of the following three conditions: BMI > 27 kg/m², insulin levels above the 90th percentile of the normal population, dyslipoproteinemia) had higher aPAF levels than those without (0.222 ± 0.167 versus 0.169 ± 0.106; p = 0.0009).

Taken together, individuals with early cardiovascular disease, as in borderline hypertension, had 5 times higher risk of being positive for aPAF than those without.

Table I

| | NT (n=73) | BHT (n=73) | p |
|-------------------------------|---------------------|---------------------|---------------|
| Waist-hip ratio | 0,90 (±0,05) | 0,92(±0,05) | 0,022 |
| Current smokers, % | 37 | 32 | |
| Cholesterol (mmol/l) | | | |
| Plasma | 5,5 (±1,0) | 5,5 (±0,9) | |
| HDL | 1,27 (±0,27) | 1,16 (±0,28) | 0,016 |
| Triglycerides (mmol/l) | | | |
| Plasma | 1,34 (±0,80) | 1,57 (±0,77) | 0,015 |
| VLDL | 0,85 (±0,69) | 1,0 (±0,68) | 0,029 |
| Insulin (mU/l) | 14,2 (±4,5) | 17,4 (±5,7) | 0,0004 |

Values are given as mean ± SD. Group differences were determined by Student's t-test or Mann-Whitney's U-test (skewed variables). HDL= high density lipoprotein, VLDL= very low density lipoprotein.

Table II. Antibody levels to PAF in subjects with or without BHT or metabolic syndrome.

| | NT (n=73) | BHT (n=73) | p |
|--------------------|--------------------|--------------------|-----------------|
| aPAF, OD405 | 0.144±0.109 | 0.215±0.130 | p=0.0007 |

Values are given as mean ± SD. Group differences were determined by Student's t-test. aPAF=antibody levels to platelet activating factor.

EXAMPLE 2

Method of determining the amount of antibodies to PAF (aPAF) in a serum sample.

IgG antibodies to PAF and lysoPAF were determined by an enzyme-linked immunosorbent assay (ELISA) essentially as described when phospholipid antibodies including cardiolipin are analysed (Harris 1986). Titertek 96-well polyvinylchloride microplates (Flow Laboratories, Costa Mesa, CA, USA) were coated with 50 µl/well of 50 µg/ml PAF dissolved in ethanol and allowed to dry overnight at 4°C. Blocking was accomplished with 20% ABS-PBS for two hours. 50 µl of serum samples, diluted 1:50 in 20% ABS-PBS were added to each well. Control assay were performed in the absence of PAF.

After 3 washings with PBS the plates were incubated with 50 µl/ml of alkaline phosphatase-conjugated goat anti-human IgG (Sigma A-3150) diluted 1:9000 with PBS at 37°C for 2 hours. After 3 washings, 100 µl of substrate (phosphatase substrate tablets, Sigma 104; 5 mg in 5 ml diethanolamine buffer, pH 9.8) was added. The plates were incubated in room temperature for 30 minutes and read in an ELISA Multiskan Plus spectrophotometer at 405 nm. Each determination was done in triplicate. The coefficient of variation between triplicate test was less than 5%.

REFERENCES CITED

Lemne C, Jogestrand T, de Faire U. Carotid intima-media thickness and plaque in borderline hypertension. *Stroke* 1995; 26:34-39.

Frostegård J, Lemne C, Andersson B, Kiessling R and de Faire, U. Association of serum antibodies to heat shock protein 65 with borderline hypertension. *Hypertension* 1997;29:40-44.

Frostegård J, Huang YH, Rönnelid J and Schäfer-Elinder L. PAF and oxidized LDL induce immune activation by a common mechanism. *Arteriosclerosis, Thrombosis and Vasc Biol.* 1997; 17: 963-968.

Barquinero J, Ordi-Ros J, Selva A, Perez-Peman P, Vilardell M, Khamashta M. Antibodies against platelet-activating factor in patients with antiphospholipid antibodies. *Lupus* 1994; 3:55-58.

Harris EN, Gharavi AE, Patel SP, Hughes GRV. Evaluation of the anti-cardiolipin antibody test: report of an international workshop held 4 April 1986. *Clin Exp Immunol* 1986;68:215-222.

U.S. patent 5,731,208

CLAIMS:

1. A method of diagnosing cardiovascular diseases in a human wherein the presence and/or concentration of antibodies to platelet activating factor is assessed in a sample of body fluid of said human.
2. The method of claim 1 wherein said cardiovascular disease is early atherosclerosis.
3. The method of claim 1 wherein said antibodies are measured by immunoassay.
4. The method of claim 1 wherein said antibodies are measured by an enzyme linked immunosorbent assay.
5. The method of claim 1 wherein said antibodies are measured by radioimmunoassay.
6. The method of claim 1 wherein the concentration is measured in serum prepared from a blood sample.
7. The method of claim 1 wherein the concentration is measured in plasma prepared from a blood sample.
8. The method of claim 1 wherein a concentration of the mean value of controls plus one standard deviation, or more, identifies an individual at high risk of having cardiovascular diseases and/or developing atherosclerosis.
9. A kit for carrying out the method according to claim 1 or 2, said kit comprising, means for testing concentration of antibodies to PAF in a sample of body fluid of a patient to be tested for cardiovascular disease and/or early atherosclerosis; and means for determining from the assayed concentration of antibodies to PAF whether the patient is at risk for having cardiovascular disease and/or early atherosclerosis.
10. A kit for diagnosing cardiovascular diseases and/or early atherosclerosis that utilizes several diagnostic indicators for said diseases, wherein antibodies to PAF is one of them.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 99/01208

A. CLASSIFICATION OF SUBJECT MATTER

IPC6: G01N 33/53, G01N 33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC6: G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category* | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|-----------|--|-----------------------|
| X | Lupus, Volume 3, 1994, Jordi Barquinero et al, "Antibodies Against Platelet-Activating Factor in Patients with Antiphospholipid Antibodies", page 55 - page 58, see page 56 | 9 |
| A | -- | 1-8,10 |
| A | Dialog Information Services, File 154, MEDLINE, Dialog accession no. 08672149, Medline accession no. 96350619, Hirashima Y et al: "Platelet-activating factor (PAF) and the formation of chronic subdural haematoma. Measurement of plasma PAF levels and anti-PAF immunoglobulin titers"; & Acta Neurochir, 1995, 137 (1-2) p15-8 | 1-10 |
| | -- ----- | |

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

5 October 1999

Date of mailing of the international search report

28.10.1999

Name and mailing address of the ISA/

Swedish Patent Office

Box 5055, S-102 42 STOCKHOLM

Facsimile No. +46 8 666 02 86

Authorized officer

Carl-Oflof Gustafsson/Eö

Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE 99/01208

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-8
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest:

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE 99/01208

Claims 1-8 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/Rule 39.1. (iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.